

Gravitational Waves from Braneworld Inflation

Andrei Frolov and Lev Kofman
CITA, University of Toronto
Toronto, Ontario, Canada, M5S 3H8
E-mail: frolov,kofman@cita.utoronto.ca
(September 2002)

We investigate the generation of primordial gravitational waves from inflation in braneworld cosmologies with extra dimensions. Advantage of using primordial gravitational waves to probe extra dimensions is that their theory depends only on the geometry, not on the microscopic models of inflation and stabilization. $D(D-3)/2$ degrees of freedom of the free bulk gravitons are projected onto the 3d brane as tensor, vector and scalar modes. We found the following no-go results for a generic geometry of a five (or D) dimensional warped metric with four dimensional de Sitter (inflationary) slices and two (or one) edge of the world branes: Massive KK graviton modes are not generated from inflation (with the Hubble parameter H) due to the gap in the KK spectrum; the universal lower bound on the gap is $\sqrt{3/2}H$. Massless scalar and vector projections of the bulk gravitons are absent, unlike in geometries with KK compactification.

A massless 4d tensor mode is generated from inflation with the amplitude H/M_P , where M_P is the effective Planck mass during inflation, derived from the D dimensional fundamental mass M_S and the volume of the inner dimensions. However, M_P for a curved dS braneworld may differ from that of the flat brane at low energies, either due to the H -dependence of the inner space volume or variations in the brane separation before stabilization. Thus the amplitude of gravitational waves from inflation in braneworld cosmology may be different from that predicted by inflation in 4d theory.

PACS numbers: 04.50.+h, 98.80.Cq

CITA-2002-22

I. INTRODUCTION

Higher dimensional formulation of superstring/M theory, supergravity and phenomenological models of the mass hierarchy has most obvious relevance to cosmology. Early universe inflation can universally amplify vacuum fluctuations of all light (minimally coupled to gravity) degrees of freedom. Thus inflation serves as a powerful tool to probe the fields associated with the extra dimensions.

In this paper we consider a class of models with extra dimensions related to the braneworld scenarios. One of the most interesting recent developments of the high energy physics phenomenology is the picture of the brane worlds, where our 3+1 dimensional spacetime is the 3d brane embedded in the higher dimensional space. In application to the very early universe this leads to the brane world cosmology, where our 3+1 dimensional universe is the 3d curved brane embedded in the bulk. A lot of efforts have been devoted to the search for the new features of the background cosmological expansion, inflation in the brane world settings, cosmological fluctuations which could probe the brane world scenarios and others. Many of these problem are rather complicated. For instance, the problem of gravitational waves which might be observed on the brane requires treatment of all $\frac{1}{2}D(D-3)$ degrees of freedom of the bulk gravitons, the choice of their initial state, the issue of the KK modes, transplanckian problem and others.

Cosmological gravitational waves are a very useful tool to probe the very early universe, because their generation

depends on the background geometry but not on the details of the particle physics built in the model. A historic example is the theory of cosmological gravitational waves generated from the early de Sitter stage [1] which, ironically, was suggested even before the microscopic models of inflation were proposed.

Gravitons almost freely propagate through the expanding universe, and may leave their prints on the power spectrum and polarization of the cosmic microwave background. With the ever increasing accuracy of the CMB anisotropy experiments, it is worth to enlarge the pool of the primordial gravitational waves theories.

In the familiar model of inflation in the four dimensions, gravitational waves are generated from vacuum fluctuations during inflation. Gravitons have two polarizations, and temporal part of their wave function $h_k(t)$ obeys an equation which is similar to the one for the massless scalar field minimally coupled to gravity.

Late time asymptotics of $h_k(t)$, which is relevant to present cosmological observations, contains long wavelength modes k . They have (almost) scale free spectrum $k^3|h_k|^2 \approx \text{const}$. The dimensionless almost constant amplitude

$$k^{3/2}h_k = \frac{H}{M_P}, \quad (1)$$

is defined by the ratio of the only two scales in the problem, which are related to the gravitational waves: the Hubble parameter during inflation H and the Planck mass M_P at low energies. Usually the Hubble parameter

slightly varies during inflation, $H = H(t)$, which brings slight tilt to the spectrum of gravitational waves (in the *r.h.s.* of (1) $H \rightarrow H(k)$). In this paper we will assume $H = \text{const}$, but will devote some discussion to the case when H slightly varies.

In this paper we will study the problem of the primordial gravitational waves generation from the early universe in the theories with large extra dimensions. As opposed to excitations of some other fields, gravitational waves propagate in the bulk. What information about the bulk/brane geometry can be, in principle, observable on the 3d brane through the gravitational excitations? The problem has the flavor of the well known mathematical quest of “how one can hear the shape of a drum?” [2].

To make geometrical setting definite, we assume the brane world inflation. We will consider geometry of five dimensional warped metric with four dimensional de Sitter slices

$$ds^2 = dy^2 + A^2(y) ds_4^2. \quad (2)$$

The warp factor $A(y)$ is determined self-consistently by the five dimensional Einstein equations, supplemented by the brane boundary conditions. This formalism is briefly outlined in Section II. For the sake of generality, we also present results for a more general case of D dimensional warped metric with $D - 1$ dimensional de Sitter slices. The class of metrics (2) covers many important brane world scenarios [3] including the Horawa-Witten theory, HW with curved branes, the Randall-Sundrum (RS) models, RSI model with phenomenological stabilization of two branes, scalar sector of the supergravity realization of the RS model and others. The presence of a bulk scalar field ϕ (a dilaton) with the potential $V(\phi)$ or several scalars is often assumed.

The warped metric (2) can be rewritten in “conformally-factorized” form

$$ds^2 = \Omega^{-2} (d\chi^2 + ds_4^2), \quad (3)$$

where conformal factor and conformal fifth coordinate are defined by

$$\Omega(\chi) = \frac{1}{A(y(\chi))}, \quad \chi = \int \frac{dy}{A(y)}. \quad (4)$$

In our treatment of gravitational waves, beginning in Section III, we will extensively use conformal mapping of the whole geometry (3), bulk and branes on its edges, into a simpler manifold with boundaries, with the metric

$$d\hat{s}^2 = \Omega^2 ds^2 = d\chi^2 + ds_4^2. \quad (5)$$

There are several important papers on the formalism of metric fluctuations [4] and properties of the gravitational waves in the brane world scenarios [5–7], and will try to make connection with them in the course of the paper. We will work with the metric in the form (3), which

allows separation of variables in gravitational wave equation, as it will be shown in Section IV. Conformal mapping also significantly simplifies the issue of the boundary conditions, as it will be explained in Section V. Just as it was in the four dimensional case, in D dimensional theory the graviton eigenfunctions $h_{k,m}(\chi)$ obey an equation for the massless scalar minimally coupled to gravity.

In Section VI, we show that the spectrum of nonzero KK modes has a mass gap with the universal lower bound $\sqrt{\frac{3}{2}} H$, independently on the bulk bulk scalar fields potential $V(\phi)$ and the presence or absence of the second brane. This result generalizes the finding of the $\frac{3}{2} H$ gap in the Randall-Sundrum II model with a single brane in the AdS bulk [6].

In Section VII, we show that the homogeneous zero KK mode $m = 0$ is degenerate, and can be described by only two physical polarization degrees of freedom, while other three can be gauged by zero by the residual coordinate transformation. Thus, the issue of the gravi-scalar and gravi-vector modes in brane cosmology is resolved: they do not contribute.

All together this allows us to conclude that the late time asymptotic of the long-wavelength gravitational waves generated from the brane inflation is defined by the two tensor polarization of the homogeneous KK mode $m = 0$, similar to what takes place in a case of inflation in four dimensions. However, as we will discuss in Section VIII, it does not mean that gravitational waves from the brane inflation in all respects are identical to those in the four dimensional theory. Essential difference arises in their amplitudes. Indeed, in five dimensional theories there is additional scale ℓ , associated with the fifth dimension (brane separation in the HW theory, the AdS radius in the RS models, etc.) With three scales in the problem, ℓ , H and M_P , one may have a dimensionless amplitude different from (1). Indeed, for the braneworld inflation, one finds $k^{3/2} h_k = \frac{H}{M_P} F(H\ell)$, with some function F . Remarkably, the amplitude can be expressed economically in terms of the effective Planck mass for curved (de Sitter) braneworld

$$k^{3/2} h_k = \frac{H}{M_{P,\text{eff}}}. \quad (6)$$

Effective Planck mass $M_{P,\text{eff}}$ during inflation in the braneworld scenario may differ from the current value $M_P \sim 10^{19}$ GeV, therefore prediction for the gravitational wave amplitude from brane inflation may differ from that from inflation in 4d theory.

In Section IX, we will discuss implications of our results to the observational test of the ratio of tensor to scalar modes, transplanckian problem in the context of models with extra dimensions, and other related issues.

We evoke the “conformally-factorized” frame (3) mainly for treatment of the gravitational waves. However, conformal mapping of the brane worlds has merits of its own. In particular, we show that the warp geometry with branes at the edges is conformally equivalent

to the portion of the spacetime with regular end of the world edges. For the case of flat 4d slices of (3), the world edges after conformal mapping are empty.

II. BACKGROUND AND GRAVITATIONAL PERTURBATIONS

In this section we outline the general formalism for treating background geometry with fields (such as dilaton) in the bulk and on the branes, and free perturbative gravitational waves in this background. In general, there may also be scalar and vector bulk perturbations as well as the perturbation in the brane position. We will not consider them. However, we will address the question if bulk gravitational perturbations generate scalar and vector perturbations at the brane.

The brane worlds scenario with Einstein gravity and scalar fields is described by the action

$$S = \frac{1}{16\pi\kappa_D^2} \int \sqrt{-g} d^D x (R + 2\mathcal{L}_{\text{bulk}}) - \frac{1}{16\pi\kappa_D^2} \sum_a \int \sqrt{-h} d^{D-1}x (2[\mathcal{K}] + 2\mathcal{L}_{\text{brane}}), \quad (7)$$

where the first term describes the bulk and the second term is related to the brane(s). The extrinsic curvature of the brane $\mathcal{K}_{\mu\nu}$ is defined through the normal unit vector n_A and the tangent vierbine $e_{(a)}^B$

$$\mathcal{K}_{\mu\nu} = e_{(\mu}^A e_{\nu)}^B \nabla_A n_B, \quad (8)$$

and $\mathcal{K}$ is its trace. We use “mostly positive” signature and curvature conventions of Misner, Thorne and Wheeler, and denote bulk indices by capital Latin letters $A, B, \dots$, brane indices by small Greek letters $\mu, \nu, \dots$, and brane hypersurfaces by Σ_a , $a = 1, 2$. We use square brackets (e.g. $[\mathcal{K}] = \mathcal{K}^+ - \mathcal{K}^-$) to denote jump of the geometrical quantity across the brane.

Einstein equations in the bulk are

$$R_{AB} - \frac{1}{2}Rg_{AB} = T_{AB}, \quad (9)$$

where the stress-energy tensor of the matter fields T_{AB} is described by the bulk Lagrangian $\mathcal{L}_{\text{bulk}}$. In particular case of the minimally coupled scalar field with potential, we take $\mathcal{L}_{\text{bulk}} = -\frac{1}{2}(\nabla\phi)^2 - V(\phi)$ and $\mathcal{L}_{\text{brane}} = U(\phi)$, and the bulk stress-energy tensor is

$$T_{AB} = \phi_{,A}\phi_{,B} - \left(\frac{1}{2}(\nabla\phi)^2 + V(\phi)\right)g_{AB}. \quad (10)$$

For 5d warped geometry (2) with scalar field the bulk Einstein equations (9) can be written as

$$\partial_y(A^{-1}\partial_y A) = -\frac{1}{4}(\partial_y\phi)^2 - \frac{1}{6}V(\phi) - (A^{-1}\partial_y A)^2, \quad (11)$$

$$6(A^{-1}\partial_y A)^2 = \frac{1}{2}(\partial_y\phi)^2 - V(\phi) + 6A^{-2}H^2, \quad (12)$$

where the second equation is a constraint. Comprehensive study of warp geometry with a scalar field based on the qualitative analysis of the dynamical system (11)-(12) was given in [10].

The jump of extrinsic curvature across the brane surface is related to the stress-energy tensor $S_{\mu\nu}$ of the matter contained on the brane by Israel’s junction conditions, which in the case of scalar field with brane potential $U(\phi)$ are

$$[\mathcal{K}_{\mu\nu} - \mathcal{K}g_{\mu\nu}] = U(\phi)g_{\mu\nu}, \quad [n \cdot \nabla\phi] = \frac{\partial U}{\partial\phi}. \quad (13)$$

From here, the assumption of Z_2 symmetry around brane Σ and the warp geometry (2) implies the boundary condition for the warp factor and the scalar field

$$\frac{\partial_y A}{A} \Big|_{\Sigma} = -\frac{U}{2(D-2)}, \quad \partial_y\phi|_{\Sigma} = \frac{1}{2} \frac{\partial U}{\partial\phi}. \quad (14)$$

In case of the RS models we have $U(\phi) = \pm\lambda$, where the sign is different for the positive or negative tension $\lambda = \text{const}$ on the brane. The bulk potential is just 5d negative cosmological constant $V(\phi) = -\Lambda$. In case of the HW model, $U(\phi) = e^{-\phi}$, $V(\phi) = e^{-2\phi}$. Our calculations are valid for arbitrary $U(\phi)$ and $V(\phi)$. As we will see, the results of the paper can be generalized for a more general case of several bulk scalars.

Free gravitational waves obey the linearized Einstein equations

$$\delta R_B^A = 0. \quad (15)$$

Linearized gravitational waves are described by the metric perturbations $h_{AB} \equiv \delta g_{AB}$, and the gravitational wave equation is

$$\square h_{AB} - 2\nabla^C \nabla_{(A} h_{B)C} + \nabla_A \nabla_B h + 2R_{(A}^C h_{B)C} = 0, \quad (16)$$

where h is a trace $h = g^{AB}h_{AB}$. In D dimensions, gravitational waves have $\frac{1}{2}D(D-3)$ independent physical degrees of freedom. In discussions of brane world perturbations, gravitational waves are often described by two components of the tensor modes, associated with the brane isometry group. However, they correspond only to the part of the bulk gravitational waves which have 5 degrees of freedom. In the formalisms of [6,7] based on the brane isometry group, one has to consider separately the pieces from scalar and vector modes to complete the description of the 5d gravitational waves.

In this paper we will deal with all five bulk graviton polarizations. We will use the “conformally-factorized” form of the background metric (3), and impose the usual transverse-traceless gauge for gravitational waves

$$\nabla^A h_{AB} = 0, \quad h = 0. \quad (17)$$

For this metric, it is convenient to fix some of coordinate freedom assuming the following orthogonality condition

$$\nabla^A \Omega h_{AB} = 0. \quad (18)$$

Similar coordinate gauge has been used in 4d cosmology where $\Omega = \Omega(t)$ (the synchronous gauge). While the two conditions (17) and (18) are not necessary compatible for a most general background, they are compatible for the brane-world metrics (2), since the orthogonality requirement is with respect to a vector $u^A = \partial/\partial\chi$ that is covariantly constant in the conformal frame $\hat{\nabla}_A u^A = 0$.

In the braneworld scenarios it is convenient to use the TT gauge (17), but one also can work with gravitational waves in other gauges. It is interesting to follow the bulk gravitational waves in the gauge which is usually used for the Kaluza-Klein compactification (however, without S^1 symmetry in the compact y direction). In the KK gauge, bulk gravitons are manifestly projected into four-dimensional world as four dimensional TT tensor mode plus scalar and vector modes. Consider the massless (homogeneous with respect to y) free scalar and vector four dimensional components. After substituting perturbations into the junction conditions (13) we derive that all light scalar and vector projection of the free bulk gravitational waves vanishes. This result will be confirmed further by calculations in the gauge (17) which we adopt in the rest of the paper.

III. GRAVITATIONAL WAVES AND CONFORMAL MAPPING

There is a useful analogy between gravitation waves in the the braneworld metric (3), and those in the 4d cosmology with the metric

$$ds^2 = a^2(\tau) (-d\tau^2 + ds_3^2), \quad (19)$$

where $ds_3 = \gamma_{ij} dx^i dx^j$ is the 3d metrics of constant curvature $K = 0, \pm 1$, $a(\tau)$ is the scalar factor of the universe and τ is a conformal time. It is well known [8] that the tensor perturbations in this metrics can be described by h_{ij} which can be conformally transformed and decomposed as $h_{ij} = \frac{1}{a} h_k(\tau) Q_{ij}$, where Q_{ij} are two-component transverse-traceless eigenmodes of ds_3^2 , $(\nabla^2 + k^2) Q_{ij} = 0$. Temporal part of the eigenmodes obeys an equation similar to that of the minimally coupled (nonconformal) scalar field

$$\ddot{h}_k + \left(k^2 - \frac{\ddot{a}}{a} + 2K \right) h_k = 0, \quad (20)$$

where dot is derivative with respect to τ .

The warped metric (3) is a higher dimensional version of (19) with time and one of the spatial coordinates exchanged. Therefore we expect that gravitational wave equation (16) can be solved in the form $h_{\mu\nu} \propto h_m(\chi) Q_{\mu\nu}$, where five gravitational polarizations are described by the transverse-traceless eigenmodes $Q_{\mu\nu}$ of 4d de Sitter spacetime, and $h_m(\chi)$ obeys an equation for minimally

coupled bulk scalar depending on conformal fifth coordinate χ . Additionally, instead of the Cauchy problem for graviton wavefunction $h_k(\tau)$ in the metric (19) we have Neumann boundary problem for graviton wavefunction $h_m(\chi)$ in the metric (3).

Let us begin with conformal transformation. All geometrical quantities and operators evaluated with respect to the original metric g_{AB} will be hatless, while those evaluated with respect to the conformally transformed metric $\hat{g}_{AB} = \Omega^2 g_{AB}$ will be denoted by hat. Covariant derivative in physical frame will be $\nabla_{\hat{\cdot}}$, while covariant derivative with respect to $\hat{g}_{AB}$ will be $\hat{\nabla}$. Under general conformal mapping, background Ricci tensor transforms as

$$R_{AB} = \hat{R}_{AB} + (D-2)\Omega^{-1}\hat{\nabla}_A\hat{\nabla}_B\Omega + \Omega^{-1}(\hat{\square}\Omega)\hat{g}_{AB} - (D-1)\Omega^{-2}(\hat{\nabla}\Omega)^2\hat{g}_{AB}, \quad (21)$$

while its trace transforms as

$$R = \Omega^2 \hat{R} + 2(D-1)\Omega(\hat{\square}\Omega) - D(D-1)(\hat{\nabla}\Omega)^2. \quad (22)$$

For gravitational perturbations we have

$$\hat{h}_{AB} = \Omega^2 h_{AB}. \quad (23)$$

It can be shown that the gauge choice (17) and (18) is conformally invariant. From the wave equation in the physical frame (16), we obtain the gravitational wave equation in the conformal frame

$$\hat{\square}\hat{h}_{AB} - 2\hat{\nabla}^C\hat{\nabla}_{(A}\hat{h}_{B)C} + 2\hat{R}^C_{(A}\hat{h}_{B)C} - (D-2)\Omega^{-1}\Omega^{;C}\hat{\nabla}_C\hat{h}_{AB} = 0, \quad (24)$$

where no assumptions about the background spacetime were made yet except the gauge (17) and (18). Although equation in the conformal frame appears to have additional term compared to (16), actually considerable simplification occurs due to the simpler form of the background metric in the conformal frame.

So far conformal transformation of the metric has been kept rather general. Now we apply the result to the conformal mapping (5) with $\Omega = \Omega(\chi)$. The curvature of the metric $d\hat{s}^2$ is determined by the curvature of the de Sitter portion, $\hat{R}_{\mu\nu} = (D-2)H^2 \hat{g}_{\mu\nu}$. Then equation (24) simplifies to

$$\hat{\square}\hat{h}_{AB} - 2\hat{\nabla}^C\hat{\nabla}_{(A}\hat{h}_{B)C} + 2(D-2)H^2\hat{h}_{AB} - (D-2)\Omega^{-1}\Omega'\hat{h}'_{AB} = 0, \quad (25)$$

where prime denotes the derivative with respect to the extra dimension χ .

IV. SEPARATION OF VARIABLES IN THE GRAVITATIONAL WAVES EQUATION

In this section, we continue the analysis of the gravitational wave equation (25). Since from the gauge (18)

it follows that $\hat{h}_{A\chi} = 0$, only $\hat{h}_{\mu\nu}$ components need to be considered. As $u^\mu = \partial/\partial\chi$ is a Killing vector for the metric in the conformal frame, the dependence on the extra dimension separates, and the gravitational perturbation can be decomposed as

$$\hat{h}_{\mu\nu} = \Omega^{\frac{D-2}{2}} h_m(\chi) Q_{\mu\nu}^{(m)}(x^\rho), \quad (26)$$

where m is an eigenvalue corresponding to the Kaluza-Klein modes. The transverse-traceless conditions (17) are inherited by the $(D-1)$ -dimensional tensor mode

$$\hat{\nabla}^\mu Q_{\mu\nu}^{(m)} = 0, \quad Q_\mu^{(m)\mu} = 0, \quad (27)$$

while no gauge conditions are placed on extra-dimensional mode h_m .

Upon substitution of the mode decomposition (26) into the equation (25), the gravitational perturbation problem is reduced to two decoupled differential equations. One of them is equation for the $(D-1)$ -dimensional tensor mode

$$\hat{\square} Q_{\mu\nu}^{(m)} - 2\hat{\nabla}^\rho \hat{\nabla}_{(\mu} Q_{\nu)\rho}^{(m)} + 2(D-2)H^2 Q_{\mu\nu}^{(m)} = m^2 Q_{\mu\nu}^{(m)}, \quad (28)$$

and the other is an equation for the extra-dimensional scalar mode

$$h_m'' + (m^2 - V_{\text{eff}}) h_m = 0, \quad (29)$$

where effective potential V_{eff} is determined by the bulk geometry

$$V_{\text{eff}} = -\frac{D-2}{2} \left(\frac{\Omega''}{\Omega} - \frac{D}{2} \frac{\Omega'^2}{\Omega^2} \right). \quad (30)$$

There is some ambiguity in the choice of the separation constant m^2 . For instance, one can include the curvature of the 4d slices into m^2 , as it is usually done for the gravitational waves in open or closed cosmology as in equation (20). Our choice of m^2 is motivated by the following reason: it turns out, as we will soon see, that m^2 in (28) and (29) has the lowest eigenvalue value $m^2 = 0$ and is always positive.

The equation (28) is *exactly* the equation for linear massive tensor perturbation of $(D-1)$ -dimensional subspace. Thus, the problem of gravitational waves in the braneworld metric (3) is reduced to two decoupled equations, one for the massive tensor polarizations $Q_{\mu\nu}^{(m)}$ on the $(D-1)$ spacetime of constant curvature which we will analyze in detail in Section VII, and one for the minimally coupled scalar mode $h_m(\chi)$. These equations are valid for a general case of an arbitrary warp factor $\Omega(\chi)$, and for flat or curved branes of constant curvature.

The equation (29) has a convenient ‘‘Schrödinger-type’’ structure. The effective potential (30) can be represented in different forms which are useful for different purposes. Using conformal transformation of the Ricci scalar (22)

$$\frac{R}{\Omega^2} = \hat{R} + 2(D-1) \left(\frac{\Omega''}{\Omega} - \frac{D}{2} \frac{\Omega'^2}{\Omega^2} \right), \quad (31)$$

the equation (29) can be written in the form

$$h_m'' + \left[m^2 + \xi_c \left(\frac{R}{\Omega^2} - \hat{R} \right) \right] h_m = 0, \quad (32)$$

and we have an alternative expression for the effective potential

$$V_{\text{eff}} = \left(\frac{D-2}{2} \right)^2 H^2 - \xi_c \frac{R}{\Omega^2}. \quad (33)$$

Numerical factor $\xi_c = \frac{1}{4} \frac{D-2}{D-1}$ corresponds to the value of the coupling constant ξ_c for a scalar field conformally coupled with gravity. This is what one expects to obtain for the equation of the minimally coupled (nonconformal) scalar field after conformal transformation. Extra-dimensional graviton mode in the physical frame corresponds to the minimally coupled massless scalar field.

If we return to the warp factor $A(\chi) = 1/\Omega(\chi)$ instead of $\Omega(\chi)$ in (30), master equation (29) will be written as

$$h_m'' + \left[m^2 - \frac{D-2}{2} \left(\frac{A''}{A} + \frac{D-4}{2} \frac{A'^2}{A^2} \right) \right] h_m = 0. \quad (34)$$

This equation generalizes equation for the 4d cosmological gravitational waves (20) (with the shifted choice of the separation eigenvalue $m^2 \rightarrow k^2 + 2K$).

Let us show that there is no tachyonic modes in the spectrum of KK modes, i.e. m^2 is always positive. The equation (29) with the potential in the form (30) can also be written in a factorized form

$$-\mathcal{D}_- \mathcal{D}_+ h_m = m^2 h_m, \quad (35)$$

where the differential operators are

$$\mathcal{D}_\pm = \partial_\chi \pm \frac{D-2}{2} (\ln \Omega)'. \quad (36)$$

Simple manipulations with the equation (35) using integration $\int d\chi$ between two branes, lead to the formula

$$m^2 = \frac{\int |\mathcal{D}_+ h_m|^2 d\chi}{\int |h_m|^2 d\chi} \geq 0, \quad (37)$$

provided that boundary terms satisfy $(h_m^* \mathcal{D}_+ h_m)|_{\Sigma_a} = 0$. This condition is met for Z_2 symmetry.

V. BOUNDARY CONDITIONS AND CONFORMAL MAPPING

The mass spectrum of the Kaluza-Klein modes is determined by a boundary value problem in the extra dimension, and is obtained by imposing desired boundary conditions on solutions of the equation (32). In this section

we discuss how the background and perturbative quantities related to boundaries are mapped under conformal transformation. The goal is to derive the boundary condition for the mode function $h_m(\chi)$. If Z_2 reflection symmetry is assumed across the brane, the constant tension brane is mapped to the edge of the conformal space $\hat{g}_{AB}$. Remarkably, it turns out that after conformal mapping the edge *regular*, i.e. is not marked by the jumps in the curvature. This conformal mapping is sketched at Figure 1.




FIG. 1. Conformal transformation simplifies geometry. Upper panel represents warped geometry (2) with branes carrying tensions on the edges. Lower panel represents conformally-factorized geometry (5) with regular edges.

The boundary condition for the gravitational wave $\hat{h}_{AB}$ in the conformal frame at the empty Z_2 -symmetric brane is simply $\hat{h}'_{\mu\nu} = 0$. Therefore, for the gravitational wave amplitude h_m , we have a simple mirror-like (Neumann) boundary conditions

$$\left(\Omega^{\frac{D-2}{2}} h_m \right)' \Big|_{\Sigma} = 0. \quad (38)$$

In the rest of this section we shall demonstrate that conformal mapping of the whole geometry, the conformally-factorized bulk metric (3) and its brane boundaries, brings it to the bulk metric (5) with empty boundaries. This brane-world conformal mapping property works for a broad class of warp geometries with the bulk scalars with flat or de Sitter branes.

Let us consider the effect of the conformal transformation $d\hat{s}^2 = \Omega^2 ds^2$ on the surface Σ of co-dimension one embedded in the spacetime manifold, and, in particular, on the extrinsic curvature $\mathcal{K}_{\mu\nu}$. Suppose the surface to be parametrized as $x^A = x^A(\xi^\mu)$, then the holonomic basis vectors tangent to the surface Σ have the same components in both physical and conformal frames

$$e^A_{(\mu)} = \frac{\partial x^A}{\partial \xi^\mu} = \hat{e}^A_{(\mu)}. \quad (39)$$

Unit normal to the surface Σ must be scaled, however, to preserve the normalization

$$n^A = \Omega \hat{n}^A, \quad n_A = \Omega^{-1} \hat{n}_A. \quad (40)$$

The extrinsic curvature (8) of the surface Σ under conformal transformation behaves as

$$\mathcal{K}_{\mu\nu} = \Omega^{-1} \hat{\mathcal{K}}_{\mu\nu} - \Omega^{-2} (\hat{\nabla} \Omega \cdot \hat{n}) \hat{g}_{\mu\nu}, \quad (41)$$

with its trace transforming as

$$\mathcal{K} = \Omega \hat{\mathcal{K}} - (D-1) (\hat{\nabla} \Omega \cdot \hat{n}). \quad (42)$$

While tangential derivatives of the conformal factor must always be continuous across the surface for induced metric to match, the normal derivative need not be so. In particular, if the brane-world has the Z_2 symmetry across the brane, $\mathcal{K}_{\mu\nu}^- = -\mathcal{K}_{\mu\nu}^+$ and $\hat{\mathcal{K}}_{\mu\nu}^- = -\hat{\mathcal{K}}_{\mu\nu}^+$, and so the normal derivative of the conformal factor flips sign $(\hat{\nabla} \Omega \cdot \hat{n})^- = -(\hat{\nabla} \Omega \cdot \hat{n})^+$. In geometrical terms, this means that the gradient of the conformal factor $\hat{\nabla} \Omega$ experiences mirror reflection on the brane.

Now let us consider what happens to junction conditions (13) under conformal transformation. The left hand side transforms as

$$[\mathcal{K}_\nu^\mu - \mathcal{K} \delta_\nu^\mu] = \Omega [\hat{\mathcal{K}}_\nu^\mu - \hat{\mathcal{K}} \delta_\nu^\mu] + (D-2) [\hat{\nabla} \Omega \cdot \hat{n}] \delta_\nu^\mu. \quad (43)$$

In the physical frame we have $[\mathcal{K}_\nu^\mu - \mathcal{K} \delta_\nu^\mu] = U(\phi) \delta_\nu^\mu$, where $U(\phi)$ is the scalar field potential at the brane. On the other hand, the last term in equation (43) gives us $(D-2) [\hat{\nabla} \Omega \cdot \hat{n}] = 2(D-2) \partial_\chi \Omega|_{\Sigma} = -2(D-2) \frac{\partial_A \Omega}{A}|_{\Sigma}$, where A is the warp factor. As can be seen from the junction condition in the form (14), these two terms in (43) are identical, and cancel. Therefore for the junction condition in the conformal frame we simply have

$$[\hat{\mathcal{K}}_\nu^\mu] = 0. \quad (44)$$

This means that in the conformal frame the brane becomes a *regular surface*, which means no special junction conditions or boundary terms are needed. The flat branes with $H = 0$ in the conformal frame becomes an empty regular surfaces.

VI. SPECTRUM OF KK MODES

Equipped with the extra-dimensional mode equation (29) and its boundary conditions (38), we can analyze its solution $h_m(\chi)$, in particular, the spectrum of KK modes for the brane world warped geometries (3).

First we notice the presence of zero mode $m^2 = 0$ homogeneous KK solution, which immediately follows from the equation in the form (35)

$$h_m(\chi) = \Omega^{-\frac{D-2}{2}}, \quad m^2 = 0. \quad (45)$$

For non-zero KK modes we have to study the eigenvalues of the Schrödinger-type equation (33), which depend on the shape of the effective potential $V_{\text{eff}}(\chi)$. Figure 2 illustrates this function for two examples of the braneworld models. Left panel corresponds to the Randall-Sundrum models (with one or two branes) with negative cosmological constant Λ in 5D AdS bulk. Right panel shows V_{eff} in the warp geometry model with bulk scalar field potential $V(\phi) = V_0 e^{-2\phi}$, corresponding to the HW model.




FIG. 2. Schrödinger potentials V_{eff} (red solid line) and $\tilde{V}_{\text{eff}}$ (blue dotted line) for the RS model (left) and HW model (right) in the periodic brane-worlds between two branes. Upper horizontal (green) line correspond to the gap $\frac{3}{2}H$ in the RS model; lower horizontal (green) line correspond to the generic gap $\sqrt{\frac{3}{2}}H$.

At first glance, the spectrum of massive KK modes crucially depends on $V_{\text{eff}}(\chi)$, and consequently, on the braneworld theory. However, it turns out that in general case one encounters the gap in the spectrum of KK modes always greater or equal to $\sqrt{\frac{3}{2}}H$. Exact numerical factor in the front of H can be slightly different (but greater) in different theories.

First, let us inspect the equation (33). If the bulk curvature is negative, $R \leq 0$, then the effective potential in the equation for the KK amplitude $h_m(\chi)$ has a lower bound

$$V_{\text{eff}} \geq \left(\frac{D-2}{2}\right)^2 H^2, \quad R \leq 0, \quad (46)$$

and the mass gap $m \geq \frac{D-2}{2}H$ appears in the spectrum. This takes place in both RS models (with one or two branes) with the AdS background supported by the negative bulk cosmological constant. The presence of the gap $\frac{3}{2}H$ in the RS model with a single brane was noticed earlier in [6] and in the domain wall in AdS model in [5].

If the bulk curvature is positive, the potential V_{eff} can be smaller than $\left(\frac{D-2}{2}\right)^2 H^2$ or even become negative. To analyze the mass spectrum in this, more general, case we will use the following trick. For a given system of eigenfunctions h_m , let us introduce adjoint system $\tilde{h}_m =$

$\mathcal{D}_+ h_m$. While h_m obey the equation in the form (35), the eigenfunctions $\tilde{h}_m$ obey the adjoint equation

$$-\mathcal{D}_+ \mathcal{D}_- \tilde{h}_m = m^2 \tilde{h}_m. \quad (47)$$

In the terminology of SUSY quantum mechanics, the system $\tilde{h}_m$ is a superpartner of h_m . The essential feature of adjoint representation for us is that the set of eigenvalues m^2 is the same for both systems h_m and $\tilde{h}_m$.

Now consider equation (47). This equation again can be rewritten in the form of the Schrödinger equation

$$\tilde{h}_m'' + (m^2 - \tilde{V}_{\text{eff}}) \tilde{h}_m = 0, \quad (48)$$

where, using the Einstein equations (11,12) for the bulk scalar field, the effective potential $\tilde{V}_{\text{eff}}$ for $\tilde{h}_m$ can be written in the form

$$\tilde{V}_{\text{eff}} = \frac{D-2}{2} H^2 + \frac{A^2}{2} (\partial_y \phi)^2 + \frac{(D-4)(D-2)}{4} (\partial_y A)^2. \quad (49)$$

In case of several bulk scalars ϕ_i gradient term $(\partial_y \phi)^2$ in (49) is replaced by positive term $\sum_i (\partial_y \phi_i)^2$. This “supersymmetric” effective potential $\tilde{V}_{\text{eff}}$ for two examples, the RS and HW models, is plotted in Figure 2 alongside with V_{eff} . From (49) one can immediately see that for an arbitrary scalar field theory

$$\tilde{V}_{\text{eff}} \geq \frac{D-2}{2} H^2. \quad (50)$$

Therefore, for $D = 5$ the spectrum of KK modes has the mass gap of at least $\sqrt{\frac{3}{2}}H$.

The presence of the gap means that there will be no KK gravitational modes (except $m = 0$ mode) which will contribute to the late time cosmological gravitational waves. This is because the allowed modes are heavy, $m \geq \sqrt{\frac{3}{2}}H$ for $D = 5$, relative to the Hubble parameter during inflation. As we will argue in the Section VII, the heavy KK modes do not survive the late-time long wavelength cosmologically relevant asymptotics.

VII. FOUR-DIMENSIONAL MASSIVE TENSOR MODES

The gravitational waves from the brane-world inflation can be factorized into the Kaluza-Klein modes $h_m(\chi)$ which obey the equation (29), and $(D-1)$ dimensional tensor modes $Q_{\mu\nu}^{(m)}(x^\rho)$ which obey the equation (28). In this section we consider $(D-1)$ tensor modes. For cosmological applications we will concentrate on $D = 5$, i.e. on four-dimensional tensor modes.

De Sitter manifold can be covered by different coordinates. It is customary in the inflationary cosmology to use a spatially flat dS slicing

$$ds_{\text{dS}}^2 = -dt^2 + e^{2Ht} dx_i dx^i \quad (51)$$

$$= a^2(\tau) (-d\tau^2 + dx_i dx^i),$$

where $i = 1, 2, 3$, $a(\tau) = -1/H\tau$. These coordinates are very convenient to study scalar fluctuations or four-dimensional massless gravitons. In paper [6], they were used to study massive transverse traceless tensor modes of the isometry group R^3 of a flat three dimensional slices of (51). As we already noted, this approach is not fully consistent, since these modes have only two polarizations and do not describe the five component bulk gravitons.

To deal with the massive transverse traceless tensor modes of dS space with the bigger isometry group $SO(4, 1)$ we shall use the global de Sitter map

$$ds_{\text{dS}}^2 = -dt^2 + \frac{1}{H^2} \cosh^2(Ht) d\Omega_3^2, \quad (52)$$

$$= \frac{1}{\cos^2 \tau} (-d\tau^2 + d\Omega_3^2);$$

$$d\Omega_3^2 = d\psi^2 + \sin^2 \psi (d\theta^2 + \sin^2 \theta d\phi^2).$$

Equation (28) for the metric (52) is the equation for tensor eigenfunctions of the Laplace-Beltrami operator on de Sitter spacetime. The iteration scheme to construct tensor eigenfunctions from scalar hyper-spherical harmonics was developed in [9].

First, we will consider the massless mode $m^2 = 0$. This is very important special case corresponding to the homogeneous KK mode. In this case the equation (28) corresponds *exactly* to the equation for the usual four dimensional massless tensor mode. Tensor mode $Q_{\mu\nu}^0$ has not five but only two transverse traceless degrees of freedom, corresponding to the two polarizations of four-dimensional graviton. Technical reason for this degeneracy is that there is a residual gauge freedom of the synchronous gauge in four dimensions which can be used to zero three components of tensor perturbations. When m^2 is non-zero, the degeneracy is lifted.

In $m^2 = 0$ case solution of the equation (28) can be further factorized and written as [9,11]

$$Q_{0\nu}^0 = 0, \quad Q_{ij}^0 = \frac{N}{a(\tau)} f_{nlm}(\tau) T_{ij}^{nlm}(\psi, \theta, \phi), \quad (53)$$

$$f_{nlm}(\tau) = \frac{1}{\sqrt{2n(n+1)(n+2)}} \times ((n+2) \cos \tau - e^{-i\tau}) e^{-i(n+1)\tau}, \quad (54)$$

where N is normalization constant independent on (n, l, m) , $T_{ij}^{nlm}(\psi, \theta, \phi)$ are transverse traceless tensor harmonics of three-sphere $d\Omega_3^2$ explicitly derived in [9], which have two independent degrees of freedom.

The key point for our discussion is that in the late time limit $t \rightarrow \infty$, $\tau \rightarrow \pi/2$, the amplitude $f_{nlm}(\tau)$ freezes out at the constant value $f_{nlm}(\tau) \rightarrow \frac{1}{\sqrt{2n(n+1)(n+2)}}$. Similar treatment of massless gravitons in the planar coordinates (52) gives us frozen amplitude $f_{\vec{k}} \rightarrow \frac{1}{(2k)^{3/2}}$ where $\vec{k}$ is 3D

comoving momentum. Therefore KK zero modes are, in principle, cosmologically observable, depending on their amplitude N .

Let us turn to the massive KK modes with $m^2 > 0$. In this case there is no degeneracy of bulk graviton degrees of freedom, and simple factorization (53) is not applicable. Massive tensor KK modes in five dimensions are described by all five components. As we shown in the previous section, in the brane inflation scenario massive gravitons are heavy, with the mass value which exceeds the universal threshold $\sqrt{\frac{3}{2}} H$. The key point about heavy gravitons is that they are not generated during inflation.

This is similar to the properties of massive scalar field fluctuations during inflation. Indeed, familiar scalar eigenfunctions of the Laplace-Beltrami operator in the de Sitter geometry in the planar coordinates are given in terms of the Hankel functions

$$f_k^{(m)}(\tau) = \frac{\sqrt{\pi}}{2} H |\tau|^{3/2} \mathcal{H}_\lambda^{(1)}(k\tau), \quad (55)$$

with the index

$$\lambda = \sqrt{\frac{9}{4} - \frac{m^2}{H^2}}. \quad (56)$$

The physical wavelength of the quantum fluctuations is stretched out with time as $\frac{a(\tau)}{k}$. Cosmologically interesting fluctuations have very small comoving momenta $k \rightarrow 0$. The amplitude of interest corresponds to the late-time asymptotics of conformal time $\tau \rightarrow 0$ (or $t \rightarrow \infty$). The late-time long-wavelengths asymptotic $k\tau \rightarrow 0$ of $f_k^{(m)}(\tau)$ crucially depends on the value of parameter $\frac{m^2}{H^2}$. For massless or very light modes, $\frac{m^2}{H^2} \ll 1$, the fluctuations are frozen out at the level

$$f_k^{(m)}(\tau) \simeq \frac{H}{\sqrt{2} k^{3/2}} \quad (57)$$

as far as its physical wavelength exceeds the Hubble radius $\frac{a(\tau_k)}{k} > H^{-1}$. On the contrary, the heavy modes $m^2 \geq H^2$ do not freeze out but oscillate with decreasing amplitude

$$f_k^{(m)}(\tau) \propto \tau^i \sqrt{\frac{m^2}{H^2} - \frac{9}{4} + \frac{3}{2}} \quad (58)$$

as $\tau \rightarrow 0$.

Therefore we conclude that *the heavy KK modes do not contribute to the cosmologically interesting late-time long-wavelengths asymptotics of the gravitational waves and scalar fluctuations*. This conclusion is similar to the result for the scalar field fluctuations in the model with inflation in outer space in theories with extra dimensions (anisotropic inflation) with the KK compactification [12]. There it was shown that heavy KK modes do not contribute to cosmological fluctuations generated from inflation.

VIII. BRANEWORLD DIMENSIONAL REDUCTION AND NORMALIZATION OF THE GRAVITATIONAL WAVE

Einstein gravitational action in D dimensions is

$$S_D = M_S^{D-2} \int R[g_{AB}] \sqrt{-g} d^D x \quad (59)$$

plus surface terms, where M_S is D -dimensional fundamental mass. If we can factorize the metric g_{AB} into four dimensional outer space and $(D - 4)$ dimensional inner space of volume V_{inner} , then the action (59) will be reduced to the effective four-dimensional Einstein action

$$S_4 = M_{P,\text{eff}}^2 \int R[g_{\mu\nu}] \sqrt{-g} d^4 x, \quad (60)$$

where the effective four-dimensional Planck mass $M_{P,\text{eff}}$ is related to the fundamental mass M_S via formula

$$M_{P,\text{eff}}^2 = M_S^{D-2} V_{\text{inner}}. \quad (61)$$

Let us follow how this procedure works for particular example of conformally-factorizable 4 + 1 metric (2) of the braneworld scenario. First we perform conformal mapping (22) and assume $\hat{R}$ does not depend on the fifth dimension χ . By doing this, we restrict ourselves to zero-mode (with respect to χ) gravitational perturbations, which is the subject of cosmological interest. By integrating extra-dimensional dependence out, we have

$$S_D = M_S^{D-2} \int \Omega^{2-D}(\chi) d\chi \int \hat{R}[\hat{g}_{AB}] \sqrt{-\hat{g}} d^{D-1} x + \dots, \quad (62)$$

where we have omitted derivative of Ω terms for brevity. The effective action for the four-dimensional theory is then

$$S_4 = M_{P,\text{eff}}^2 \int R[g_{\mu\nu}] \sqrt{-g} d^4 x + \dots, \quad (63)$$

where effective four-dimensional Planck mass is

$$M_{P,\text{eff}}^2 = M_S^3 \ell_{\text{eff}} \quad (64)$$

and ℓ_{eff} is the effective size (volume) of the inner dimension. For braneworld scenario with compact inner dimension χ , it is

$$\ell_{\text{inn}} = \int_{\chi_1}^{\chi_2} \Omega^{-3}(\chi) d\chi = \int_{y_1}^{y_2} A^2(y) dy. \quad (65)$$

The warp factor $A(y)$, in principle, depends on the brane curvature (defined by H) through the Einstein equation (12). In particular, it is different for de Sitter brane geometry with the Hubble parameter H and for the flat brane with $H = 0$. Independently, physical separation

between curved branes may not be the same as the separation of flat branes at low energies. Indeed, in the braneworld cosmology energy-momentum tensor at the brane is time dependent, and so is the extrinsic curvature, i.e. the brane embedding. Therefore the effective volume ℓ_{eff} of the extra dimension is changing in the course of cosmological evolution between inflation and present universe.

Thus, the effective Planck mass $M_{P,\text{eff}}$ should be expected to be different during inflation and in the present universe.

Let us return to the gravitational waves. In the actions (59), (60) we split the curvature into the background part plus gravitational wave perturbations. To construct the graviton field theory we keep perturbations up to the second order. Contribution to the action from the second order graviton perturbations reads as

$$\delta S_5 = M_S^3 \int \Omega^{-3} \left[\frac{1}{4} \hat{h}_{AB;C} \hat{h}^{AB;C} + \dots \right] \sqrt{-\hat{g}} d^5 x. \quad (66)$$

As we found in the previous section, only zero KK mode contributes to the interesting cosmological effects. For these modes integrand in (66) does not depend on the fifth dimension, and we obtain

$$\delta S_4 = M_{P,\text{eff}}^2 \int \left[\frac{1}{4} h_{\mu\nu;\rho} h^{\mu\nu;\rho} + \dots \right] \sqrt{-g} d^4 x. \quad (67)$$

Thus, we reduce the problem back to the four-dimensional theory of gravitational waves from inflation. The standard approach is to rewrite action (67) in terms of scalar field φ with the canonical kinetic term, $\varphi = \frac{M_{P,\text{eff}}}{\sqrt{2}} h$, and quantize the scalar field φ in the de Sitter background. Scalar field vacuum fluctuations are amplified by inflation to the magnitude $k^{3/2} \varphi_k = \frac{H}{\sqrt{2}}$ (see also (57)). As a result, we obtain that long-wavelength gravitational waves are generated from inflation with the scale free spectrum and the amplitude

$$k^{3/2} h_k = \frac{H}{M_{P,\text{eff}}}. \quad (68)$$

Thus, the only difference with the results for the gravitational wave amplitude in the usual four-dimensional theory is the effective Planck mass $M_{P,\text{eff}}$ instead the fixed M_P , namely

$$k^{3/2} h_k = \frac{H}{M_P} \sqrt{\frac{\ell_{\text{now}}}{\ell_{\text{inf}}}}, \quad (69)$$

where ℓ_{now} , ℓ_{inf} are the effective sizes of extra dimension at the inflationary stage and today.

Let us discuss how $M_{P,\text{eff}}$ at the stage of inflation in the braneworld scenario may be different from the current M_P . It is instructive to consider the simple Randall-Sundrum model with a single inflating brane in five dimensional AdS bulk. There we have the conformal factor $\Omega(\chi) = \frac{\sinh H\chi}{H\ell}$. Then explicit integration in (65) gives [6]

$$\ell_{\text{eff}} = \ell \left\{ \sqrt{1 + H^2 \ell^2} + H^2 \ell^2 \ln \left(\frac{\sqrt{1 + H^2 \ell^2} - 1}{H \ell} \right) \right\}. \quad (70)$$

For the ratio $\frac{\ell_{\text{now}}}{\ell_{\text{inf}}}$ involved in formula (69) we have the asymptotics

$$\frac{\ell_{\text{now}}}{\ell_{\text{inf}}} \rightarrow \begin{cases} 1 & \text{for } H\ell \ll 1 \\ \frac{3}{2}(H\ell) & \text{for } H\ell \gg 1 \end{cases}. \quad (71)$$

If brane inflation occurred at scales smaller than the curvature of the extra dimension (i.e. $H\ell \gg 1$), and today universe is almost flat (i.e. $H\ell \ll 1$), the gravitational wave amplitude generated by inflation is enhanced

$$k^{3/2} h_k = \frac{H}{M_{P,\text{eff}}} = \frac{H}{M_P} \sqrt{\frac{3}{2} H\ell} \quad (72)$$

compared to our expectation from standard cosmology.

Another example where $M_{P,\text{eff}}$ is changing in the braneworld scenario was considered in [7]. It can be shown that the model of [7] effectively corresponds to the case of two parallel inflating branes separated by scale ℓ embedded in flat empty five dimensional space. There it was assumed that $H\ell \ll 1$.

We are not aware of convincing arguments why $H\ell$ cannot be of order of unity or larger. The case of $H\ell \ll 1$ corresponds to two (almost) parallel branes separated by the scale $\sim \ell$ which is small compared to their curvature radius H^{-1} . It looks natural to have as initial conditions two branes parallel within the casually connected region [13]. From point of view of four dimensional observer this is the region $\sim H^{-1}$. However, casually connected region in five dimensional space is defined not by H but by five dimensional curvature. For instance, in the RSI scenario with the AdS bulk already for $H\ell \sim 1$ the function ℓ_{eff} in (70) drops by factor 2 (it is a very steep function of $H\ell$). The case $H\ell \geq 1$ gives us interesting possibility to have significant departure from the standard prediction of the inflation in the four dimensional theory.

So far we discussed the amplitude of the gravitational waves at the stage of inflation. Will formula (68) be the final answer for the gravitational wave amplitude in the late time cosmology? The reason for this question is the following. The amplitude h_k may somehow be affected during cosmological evolution between inflation and the latest stages because $M_{P,\text{eff}}$ may change its value between inflation and current value (say, after stabilization after inflation).^{*} Although equation for free gravitational waves does not depends on the gravitational coupling, situation with the varying M_P is more delicate. One may model this situation in framework of the Brans-Dicke theory, where BD scalar has two constant

but different asymptotics. Simple estimations with this approach indicates that there is no additional variation of the amplitude h_k . This issue deserves further investigation.

IX. DISCUSSION

Among the many features of superstring/M theory, the existence of extra dimensions has the most direct relevance to cosmology. Gravitons propagate in all – outer and inner – dimensions. Bulk gravitons in D dimensions have $\frac{1}{2}D(D-3)$ components. Only two of them appear as a transverse traceless tensor mode from the point of view of a four dimensional observer, while the others are projected into the 3 + 1 dimensional world as (gravi)scalars and (gravi)vectors.

Vacuum fluctuations of all light (minimally coupled) degrees of freedom are unstable during inflation and appear as long-wavelength perturbations in late time cosmology. Thus, outer space inflation acts like an amplifier and stretcher of light modes. Using inflation as a tool, we may probe extra dimensions by examining light modes associated with extra dimensions which may be produced by inflation.

It is especially appealing to investigate gravitational waves in the context of extra dimensions plus inflation. This is because the theory of small gravitational fluctuations depends on the underlying geometry but not on the microscopic model of its realization. Additionally, gravitons propagate almost freely throughout the evolution of the universe, thus carrying information about the primordial universe.

There are two distinct types of scenarios with compactification of extra dimensions from the point of view of the theory of primordial gravitational waves. One type is Kaluza-Klein compactification. The theory of gravitational waves from inflation in models with KK compactification will be considered separately. Another type of model arises from the braneworld scenarios with heavy branes (including a single brane in AdS bulk or compact warped inner space in between two branes). In this paper we investigated primordial gravitational waves in generic geometries of the warped inner space with outer space (brane) inflation.

We found the following no-go results for gravitational waves from inflation in the generic case of these braneworld scenarios:

Massless mode of bulk gravitons is projected to the usual four dimensional two component TT tensor mode, while their scalar and vector projections are absent. This is different from KK compactification where light bulk gravitons are projected to tensor, vector and scalar massless components in the four dimensional world.

Massive KK modes are not generated from inflation due to the gap in the KK spectrum. The gap is present in all models including models with two branes. The

^{*}We thank Shinji Mukohyama for this comment.

universal lower bound of the gap is

$$\Delta m \geq \sqrt{\frac{3}{2}} H. \quad (73)$$

This result generalizes the finding of the gap (of magnitude $\frac{3}{2}H$) for the RS model with a single brane embedded in AdS [6].

Among all massless projections of the bulk gravitons, only the four dimensional TT massless modes are generated from brane inflation. As in the standard four dimensional theory, their spectrum is scale free, but the amplitude is given by the formula

$$k^{3/2} h_k = \frac{H}{M_{P,\text{inf}}}, \quad (74)$$

where $M_{P,\text{inf}}$ is the four dimensional Planck mass for the curved (de Sitter) observable brane. Suppose the inner dimensions are stabilized after inflation (say due to SUSY breaking). This is not too radical an assumption bearing in mind that the usual energy scale of inflation is $H \sim 10^{13}$ GeV and the usual energy scale of SUSY breaking is $\sim 10^{11}$ GeV. Therefore theoretically it is not necessary that $M_{P,\text{inf}}$ during inflation coincides with present day low energy M_P . The difference may come from variation of the size of the inner dimension(s) (i.e. the limits of integration in formula (65), or from an imprint of the brane curvature H on the warp factor $A(y)$. A change in $M_{P,\text{eff}}$ by a factor of 2 is already interesting. Thus in general, the connection between gravitational wave amplitudes and energy scale of inflation H in theories with extra dimensions may be different from that in four dimensional theory.

Most important test of the primordial cosmological fluctuations from inflation is the ratio between the amplitude of tensor and scalar perturbations T/S . To calculate scalar perturbations, one needs to specify the model of inflation (say in terms of the scalar field potential at the brane). Already the simplest model of inflation in the RSI scenario without bulk scalars predicts that T/S may be different from that predicted in the four dimensional theory. The theory of scalar perturbations in braneworld scenarios with bulk scalars is even more involved. In general, we expect that T/S from braneworld inflation will be model-dependent and depend on many parameters.

Finally, we will make a comment on the transplanckian problem in the context of extra dimensions plus inflation. The idea beyond this problem is the following [14]: In an expanding universe the physical momentum of quantum fluctuations is redshifted to smaller values. Going backwards in time, we have to deal with ever increasing momenta which at some instant will be equal to the fundamental mass scale M_S . At this point and beyond the quantum field theory approach to fluctuations in an expanding universe (say during inflation) should be replaced by a string theory description, which is not yet available. For quantum fluctuations from inflation in the

4d theory there is, however, a convenient phenomenological coding of the problem in terms of the Bogolyubov coefficients α_k, β_k for the initial vacuum state: instead of the vacuum eigenmode $f_k(\tau)$ at infinity (given in terms of the Hankel function (55)) one can use the mode function

$$f_k(\tau) \rightarrow \alpha_k f_k(\tau) + \beta_k f_k^*(\tau), \quad |\alpha_k|^2 - |\beta_k|^2, \quad (75)$$

and discuss further the effects from β_k , if any. If the sizes of the extra dimensions are larger than M_S^{-1} which takes place in many phenomenological schemes (like the HW theory), then we must simultaneously deal with extra dimensions and the transplanckian problem. It is, however, straightforward to combine the formalism of cosmological fluctuations and that of the transplanckian problem. If the sizes of the extra dimensions are large enough to work there in the supergravity approximation, we can use the quantum field theory of fluctuations in the bulk, as we did in this paper. Then the transplanckian story will again be coded in terms of β_k for bulk quantum eigenmodes (of gravitons, for instance).

ACKNOWLEDGMENTS

We are grateful to R. Brandenberger, G. Felder, Y. Dolivet, W. Israel, A. Linde, S. Mukohyama, S. Pichler, A. Starobinsky and D. Wands for valuable discussions. This research was supported in part by the Natural Sciences and Engineering Research Council of Canada and CIAR.

-
- [1] A. A. Starobinsky, *Spectrum of relict gravitational radiation and the early state of the universe*, JETP Lett. **30**, 682 (1979) [Pisma Zh. Eksp. Teor. Fiz. **30**, 719 (1979)].
 - [2] M. Kac, *Can one hear the shape of a drum?*, Amer. Math. Monthly, 73 (1966).
 - [3] P. Horava and E. Witten, *Heterotic and type I string dynamics from eleven dimensions*, Nucl. Phys. B **460**, 506 (1996) [[hep-th/9510209](#)]; A. Lukas, B. A. Ovrut, K. S. Stelle and D. Waldram, *Heterotic M-theory in five dimensions*, Nucl. Phys. B **552**, 246 (1999) [[hep-th/9806051](#)]; L. Randall and R. Sundrum, *A large mass hierarchy from a small extra dimension*, Phys. Rev. Lett. **83**, 3370 (1999) [[hep-ph/9905221](#)]; L. Randall and R. Sundrum, *An alternative to compactification*, Phys. Rev. Lett. **83**, 4690 (1999) [[hep-th/9906064](#)]; W. D. Goldberger and M. B. Wise, *Modulus stabilization with bulk fields*, Phys. Rev. Lett. **83**, 4922 (1999) [[hep-ph/9907447](#)]; O. DeWolfe, D. Z. Freedman, S. S. Gubser and A. Karch, *Modeling the fifth dimension with scalars and gravity*, Phys. Rev. D **62**, 046008 (2000) [[hep-th/9909134](#)].
 - [4] S. Mukohyama, *Gauge-invariant gravitational perturbations of maximally symmetric spacetimes*, Phys. Rev.

- D **62**, 084015 (2000) [[hep-th/0004067](#)]; H. Kodama, A. Ishibashi and O. Seto, *Brane world cosmology: Gauge-invariant formalism for perturbation*, Phys. Rev. D **62**, 064022 (2000) [[hep-th/0004160](#)]; D. Langlois, *Brane cosmological perturbations*, Phys. Rev. D **62**, 126012 (2000) [[hep-th/0005025](#)]; C. van de Bruck, M. Dorca, R. H. Brandenberger and A. Lukas, *Cosmological perturbations in brane-world theories: Formalism*, Phys. Rev. D **62**, 123515 (2000) [[hep-th/0005032](#)]; K. Koyama and J. Soda, *Evolution of cosmological perturbations in the brane world*, Phys. Rev. D **62**, 123502 (2000) [[hep-th/0005239](#)]; N. Deruelle, T. Dolezel and J. Katz, *Perturbations of brane worlds*, Phys. Rev. D **63**, 083513 (2001) [[hep-th/0010215](#)].
- [5] S. W. Hawking, T. Hertog and H. S. Reall, *Brane new world*, Phys. Rev. D **62**, 043501 (2000) [[hep-th/0003052](#)].
- [6] D. Langlois, R. Maartens and D. Wands, *Gravitational waves from inflation on the brane*, Phys. Lett. B **489**, 259 (2000) [[hep-th/0006007](#)].
- [7] G. F. Giudice, E. W. Kolb, J. Lesgourgues and A. Riotto, *Transdimensional physics and inflation*, [[hep-ph/0207145](#)].
- [8] H. Kodama and M. Sasaki, *Cosmological perturbation theory*, Prog. Theor. Phys. Suppl. **78**, 1 (1984).
- [9] A. Higuchi, *Symmetric tensor spherical harmonics on the N -sphere and their application to the de Sitter group $SO(N, 1)$* , J. Math. Phys. **28**, 1553 (1987); A. Higuchi, *Quantum linearization instabilities of de Sitter space-time. 1*, Class. Quant. Grav. **8**, 1961 (1991); A. Higuchi, *Quantum linearization instabilities of de Sitter space-time. 2*, Class. Quant. Grav. **8**, 1983 (1991); A. Higuchi, *Linearized gravity in de Sitter space-time as a representation of $SO(4, 1)$* , Class. Quant. Grav. **8**, 2005 (1991).
- [10] G. N. Felder, A. Frolov and L. Kofman, *Warped geometry of brane worlds*, Class. Quant. Grav. **19**, 2983 (2002) [[hep-th/0112165](#)].
- [11] B. Allen, *The stochastic gravity wave background in inflationary universe models*, Phys. Rev. D **37**, 2078 (1988).
- [12] L. A. Kofman and A. A. Starobinsky, *Instability of a scalar field in a geometry with anisotropic inflation*, Phys. Lett. B **188**, 399 (1987).
- [13] N. Kaloper and A. D. Linde, *Inflation and large internal dimensions*, Phys. Rev. D **59**, 101303 (1999) [[hep-th/9811141](#)].
- [14] G. Chibisov and Yu. Shtanov, *Structural anisotropy in a chaotic inflationary universe*, Sov. Phys. JETP **69**, 17 (1989); R. H. Brandenberger and J. Martin, *The robustness of inflation to changes in super-Planck-scale physics*, Mod. Phys. Lett. A **16**, 999 (2001) [[astro-ph/0005432](#)]; A. A. Starobinsky, *Robustness of the inflationary perturbation spectrum to trans-Planckian physics*, Pisma Zh. Eksp. Teor. Fiz. **73**, 415 (2001) [JETP Lett. **73**, 371 (2001)] [[astro-ph/0104043](#)]; N. Kaloper, M. Kleban, A. E. Lawrence and S. Shenker, *Signatures of short distance physics in the cosmic microwave background*, [[hep-th/0201158](#)]; R. H. Brandenberger and J. Martin, *On signatures of short distance physics in the cosmic microwave background*, [[hep-th/0202142](#)]; R. Easther, B. R. Greene, W. H. Kinney and G. Shiu, *A generic estimate of trans-Planckian modifications to the primordial power spectrum in inflation*, Phys. Rev. D **66**, 023518 (2002) [[hep-th/0204129](#)].